



UNIVERSITI PUTRA MALAYSIA

**BIOLOGICAL CONTROL OF THE DIAMONDBACK MOTH
(PLUTELLA XYLOSTELLA LINNAEUS) USING PLUTELLA
XYLOSTELLA GRANULOVIRUS**

AHMAD DEZIANIAN

FH 2011 3

**BIOLOGICAL CONTROL OF THE DIAMONDBACK MOTH (*PLUTELLA*
XYLOSTELLA LINNAEUS) USING *PLUTELLA XYLOSTELLA*
GRANULOVIRUS**

By

AHMAD DEZIANIAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

February 2011



DEDICATION

I would like to dedicate my thesis to my late father, my dear mother and beloved spouse who always inspired and supported me to achieve this goal

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the degree of Doctor of Philosophy

**BIOLOGICAL CONTROL OF THE DIAMONDBACK MOTH (*PLUTELLA*
XYLOSTELLA LINNAEUS) USING *PLUTELLA XYLOSTELLA*
GRANULOVIRUS**

By

AHMAD DEZIANIAN

February 2011

Chairman: Professor Ahmad Said Sajap, PhD

Faculty: Forestry

The diamondback moth (DBM) *Plutella xylostella*, is known to be the most serious pest of crucifers in the world. Regular and repeated use of chemical insecticides for controlling the pest over the years has resulted in DBM developing resistance to all classes of chemical insecticides. These shortcomings have prompted researchers to evaluate the potential of biological control of DBM as an alternative to conventional insecticides. One of the biological control agents is *Plutella xylostella* granulovirus. *Plutella xylostella* granulovirus (PxGV) was effective in suppressing the Malaysian population of DBM larvae. The virus was originally isolated from Taiwan and was confirmed by Restriction Endonuclease (REN) analysis. The pattern and DNA profile of the GV was compared with that of PxGV isolates reported earlier. Scanning Electron Micrograph revealed that the capsules of PxGV were ovocylindrical with a mean size of 272.84 ± 12 (length) \times 148.27 ± 19 (width) nm. The virions were 168.44 ± 16 (length) \times 29.57 ± 12 (width) nm and rod-shaped. Results from bioassay showed that larval mortality was significantly influenced by larval instar and viral concentrations. Mortalities caused by the highest virus

concentration (3.11×10^9 Granules/ml) were 98.30 %, 96.60 %, 73.30 %, and 14.30 %, in the first to fourth instar, respectively. Larval mortality increased with increasing viral concentrations and decreased with the older instars. The median lethal concentration (LC_{50}) in the second instars was estimated 1.39×10^6 Granules/ml.

PxGV was susceptible to ultra violet radiation (UV) and lost almost all of its infectivity after 7 hours of exposure to UV-B radiation under laboratory conditions. Virulency of UV treated *PxGV* were reduced when compared to non-treated *PxGV* as much as 19.64%, 41.53%, 63.17%, 70% and 89 % after 5, 15, 30, 60 and 120 minutes exposure to UV radiation, respectively. Adjuvants consisting of Tinopal, molasses, lignin and skimmed milk added separately to *PxGV* suspension significantly improved the residual activity of *PxGV* after exposure to UV radiation. *PxGV* + Tinopal, *PxGV* + molasses, *PxGV* + lignin and *PxGV* + skimmed milk increased residual activity 67.78%, 65.31%, 59.55% and 31.35% after being exposed to UV radiation, respectively. The molasses and Tinopal when incorporated with different virus concentrations before exposure to UV light significantly increased the residual activity. Molasses+virus showed greatest effects on the larval mortality at all virus concentrations compared to those of Tinopal+virus and lignin+virus before exposure to UV light. The LC_{50} calculated for virus + molasses (5.2×10^4 Granules/ml) before exposure to UV light was 9.2 and 1.75 times lower than lignin+virus and Tinopal+virus respectively.

Four spray-dried formulations were produced using a spray drying machine (Mini Spray Dryer Buchi-B 290) at concentration of 1.4×10^{10} Granules/g and coded as

CLM (*PxGV* + modified food starch + lignin + molasses), KLM (*PxGV* + kaolin + lignin + molasses), CAP (*PxGV* + modified food starch) and KAL (*PxGV* + kaolin). These formulations, except KAL, caused higher larval mortality than the unformulated virus before exposure to UV light. The CLM formulation caused the highest mortality (98.60%) after exposure to UV light and significantly improved the residual activity compared to the unformulated virus. KLM, CAP, KAL formulations and control treatment provided 87.13%, 64.87%, 45.97% and 27.06% larval mortality, respectively. The LC_{50} values estimated for CLM formulations (1.7×10^5 Granules/ml) was 1.76 fold lower than KLM formulation. The results indicate that spray drying procedure did not cause adverse effects on the virus particles. The encapsulated formulation not only provided effective protection to occlusion bodies against UV radiation but it also improved biological activity of the virus as well. The spray-dried ingredients and size of microgranules considerably improved CLM formulation suspensibility. The results of semi-field tests showed that CLM formulation considerably improved residual activity of the *PxGV* under natural conditions. After 0.5, 5, 24, 48 and 72 hours of exposure to the natural condition, virus of the CLM formulation still retained 13.7%, 30.67%, 42.93%, 52.53% and 21.76% compared to unformulated virus, respectively. CLM formulation was significantly more effective on larval mortality than KLM formulation after exposure to natural conditions. Although spray dried formulation of *PxGV* was effective for controlling of the pest both in the laboratory and semi-field tests, further field experiment and economical feasibility study are recommended.

Abstrak of tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

PENGAWALAN BIOLOGI BAGI DIAMONDBACK MOTH (*PLUTELLA XYLOSTELLA*.LINNAEUS) DENGAN MENGGUNAKAN *PLUTELLA XYLOSTELLA* GRANULOVIRUS

Oleh

AHMAD DEZIANIAN

Febuari 2011

Pengerusi: Profesor Ahmad Said Sajap, PhD

Fakulti: Perhutanan

Diamondback moth (DBM) *Plutella xylostella* dikenali sebagai perosak yang sangat serius bagi sayur-sayuran di dunia. Penggunaan racun kimia untuk pengawalan serangga perosak yang bertahun-tahun lamanya, telah memberikan keputusan dalam peningkatan sistem pertahanan DBM kepada semua pengkelasan bagi racun kimia. Situasi ini telah mendorong para penyelidik mengambil langkah segera untuk menilai potensi pengawalan biologi bagi DBM sebagai langkah alternatif racun kimia. Salah satunya adalah *P. xylostella* granulovirus (*PxGV*) sebagai agen pengawalan biologi yang berkesan untuk mengawal populasi bagi larva DBM di Malaysia. Virus ini berasal dari Taiwan dan telah disahkan oleh analisis Restriction Endonuclease (REN). Size genome GV telah dipatenkan berbanding dengan isolate *PxGV* yang telah dilaporkan. Melalui pemerhatian Scanning Electron Micrograph, dari segi ciri morfologi, *PxGV* telah mempamerkan kapsulnya berbentuk ovocylindrical dengan bersaiz $272.84 \pm 12 \times 148.27 \pm 19$ nm. Manakala virion-virionnya adalah bersaiz $168.44 \pm 16 \times 29.57 \pm 12$ nm dan berbentuk rod. Keputusan

bioassay telah mempamerkan kematian larva adalah significant berdasarkan peringkat larva dan kepekatan viral. Kematian disebabkan kepekatan virus yang tinggi (3.11×10^9 Granules/ml) di mana 98.3%, 96.6%, 73.3%, dan 14.3 % daripada instar pertama hingga keempat masing-masing. Kematian larva meningkat dengan peningkatan kepekatan dan pengurangan bagi instar yang semakin matang. Pertengahan lethal kepekatan (LC_{50}) dalam instar kedua adalah dianggarkan 1.39×10^6 Granules/ml.

PxGV adalah sensitif dibawah ultra ungu dan kebanyakannya hilang sifat kepathogenan selepas terdedah selama tujuh jam. Kehilangan sifat kepathogenan virus masing-masing telah mencatat 19.64%, 41.5%³, 63.17%, 70% and 89 % selepas terdedah selama 5, 15, 30, 60 hingga 120 minit di bawah radiasi UV. Bahan-bahan yang meliputi Tinopal, molasses, lignin dan susu skim ditambah secara berasingan pada *P. xylostella* granulovirus (*PxGV*) adalah signifikan dengan peningkatan aktiviti bagi residual *PxGV* selepas terdedah di bawah radiasi UV. *PxGV* + Tinopal, *PxGV* + molasses, *PxGV* + lignin dan susu skim + *PxGV* meningkat 67.78%, 65.31%, 59.55% dan 31.35 % aktiviti residualnya selepas terdedah di bawah radiasi UV masing-masing. Gabungan Molasses dan Tinopal pada kepekatan virus yang berbeza sebelum terdedah di bawah sinaran UV adalah significant bagi peningkatan aktiviti residual. Molasses + virus menunjukkan kesan yang sangat memberansangkan bagi kematian larva pada kesemua kepekatan berbanding dengan Tinopal + virus dan lignin + virus sebelum terdedah di bawah sinaran cahaya UV. LC_{50} menunjukkan virus + molasses (5.2×10^4 Granules/ml) adalah 9.2 dan 1.75 kali kurang daripada lignin + virus dan Tinopal + virus masing-masing sebelum terdedah di bawah sinaran UV.

Empat formula semburan pengeringan telah dihasilkan menggunakan mesin semburan pengeringan (Mini Spray Dryer Buchi-B 290) pada kepekatan 1.4×10^{10} OB/g dan dikodkan sebagai CLM (*PxGV* + modified food starch + lignin + molasses), KLM (*PxGV* + kaolin + lignin + molasses), CAP (*PxGV* + modified food starch) dan KAL (*PxGV* + kaolin). Formula-formula tersebut kecuali KAL menunjukkan peningkatan dalam kematian larva berbanding virus tanpa formula sebelum terdedah di bawah sinaran cahaya UV. Formula CLM menunjukkan kematian yang tinggi (98.60%) selepas terdedah di bawah sinaran UV dan significant bagi peningkatan aktiviti residual berbanding dengan virus tanpa formula. Manakala formula KLM, CAP, KAL dan rawatan kawalan masing-masing menghasilkan 87.13%, 64.87%, 45.97% dan 27.06 % bagi kematian larva. Nilai LC_{50} dianggarkan untuk formula CLM adalah $(1.7 \times 10^5 \text{ Granules/ml})$ 1.76 bahagian adalah kurang daripada formula KLM. Keputusan-keputusan tersebut mendapati prosedur penyemburan kering tidak memberi sebarang kesan terhadap partikel virus. Lagipun formula encapsula bukan hanya menghasilkan perlindungan kepada jasad virus di bawah radiasi UV yang banyak tetapi juga meningkatkan aktiviti biologi yang sangat baik bagi virus. Kandungan semburan pengeringan dan saiz bagi microgranul didapati memberi peningkatan suspensibiliti bagi formula CLM. Keputusan-keputusan bagi ujian separuh lapangan menunjukkan peningkatan aktiviti residual formula CLM bagi *PxGV* di bawah keadaan semulajadi. Selepas terdedah selama 0.5, 5, 24, 48 dan 72 jam kepada keadaan semulajadi, virus bagi formula CLM masih mengekalkan 13.7%, 30.67%, 42.93%, 52.53% dan 21.76 % aktiviti residualnya berbanding dengan tanpa formula masing-masing. Formula CLM adalah significant dan sangat berkesan terhadap kematian larva berbanding formula KLM selepas terdedah pada keadaan semulajadi. Walaupun formula penyemburan kering *PxGV*

menunjukkan keberkesanan yang menarik untuk mengawal serangga perosak
baikpun di dalam makmal dan separa lapangan, eksperimen di lapangan dan
kemampuan ekonomi dicadangkan.

ACKNOWLEDGEMENTS

In the name of Allah, The Most Merciful and Most Benevolent

This dissertation would not have been finished if it were not for the help and support of following individuals and institutions.

I am extremely thankful to my main supervisor Prof. Dr. Ahmad Said Sajap for his continuous support, warm encouragement and thoughtful guidance in the Ph.D. program. I wish to express my sincere thanks to worthy members of my supervisory committee: Dr. Lau Wei Hong, whose helpful guidance and support during laboratory works, Prof. Dr. Dzolkhifli Bin Omar who gave insightful comments, Dr. Rozi Mohamed for her encouragement and comments and Dr. Mohamed Rani Mat Yusoh whose helpful suggestion. I owe my deepest gratitude to Mr. Hussan Abdul Kadir for provision of virus stocks, his friendship useful guidance and helpful discussion. I also thank Madam Siti Noor Aishikin A.H for laboratory and insectarium facilities and two other insectarium staff, Madam Fatimah Bt Harun and Miss Dayangku Rospinar Awannku Kadree for DBM rearing larvae in MARDI. I also thank to MSc. student Mr. Bong Chee Bue for his friendship assistance during virus purification and DNA isolation in Microbiological Laboratory of Agricultural Faculty.

My special thanks to Dr. David Grzywacz from Natural Resources Institute, University of Greenwich, UK for his helpful advice relating to molecular biology works, Dr. Robert Behle for his helpful guidance, Dr. Robert Farrar and Dr. Lawrence A Lacey from USDA who has never tired of answering my questions



relating to virus formulation. I am also grateful to Dr. Ahmad Selamat for assistance in analysis of data. I would like to thank Mr. Ho Oi Kuan and Madam Farah Deba staff of Institute of BioScience (IBS) at UPM (Universiti Putra Malaysia) for scanning electron microscopy (SEM) of *PxGV* and microgranules. I also thank to Mr. Ahmad Ridhwan Isa and Miss Rafizah Binti Minhat, entomology laboratory staff of Forestry Faculty for collaboration in Sawi plantation. I wish to express my special thanks to Mr. Mohamad Soib Yusof and Mr. Mohd Amran Suratman, laboratory assistants of Food Engineering Laboratory of Food Science and Technology Faculty who taught me how to use the spray drying machine and assistance during spray drying process. I am grateful to Mr. AB. Aziz Abdollah assistant science officer of Physic and Soil Conservation Laboratory of Agricultural Faculty for meteorological information.

I would also like to acknowledge Malaysian Agricultural Research and Development Institute (MARDI) for allowing me to use their laboratory facilities and greenhouse to conduct this study.

Lastly, I wish to express my gratitude to my family for love and encouragement. For my darling mother, for her prayers and moral support, for my late father who raised me with a love of science and supported me in all my pursuits. And the most of all for my loving, supportive, encouraging, and patient wife, Homa whose faithful support during the stages of this Ph.D. is so appreciated.

I certify that a Thesis Examination Committee has met onto conduct the final examination of Ahmad Dezianian on his thesis entitled “Biological Control of Diamondback Moth (*Plutella xylostella* Linnaeus) Using *Plutella xylostella* granulovirus” with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U. (A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Examiner 1, PhD
(Chairman)

Examiner 2, PhD
(Internal Examiner)

Examiner 3, PhD
(Internal Examiner)

Examiner 4, PhD
Professor
(External Examiner)

HASANAH MOHD GHAZALI, PhD
Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 19 April 2011

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Ahmad Said Sajap, PhD

Professor
Faculty of Forestry
Universiti Putra Malaysia
(Chairperson)

Dzolkhifli Omar, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Lau Wei Hong, PhD

Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Rozi Mohamed, PhD

Faculty of Forestry
Universiti Putra Malaysia
(Member)

Mohamed Rani Mat Yusoh, PhD

Malaysian Agricultural
Research and Development
Institute, MARDI
(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 19 April 2011



DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

AHMAD DEZIANIAN

Date: 22 February 2011

TABLE OF CONTENTS	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENT	x
APPROVAL	xii
DECLARATION	xiv
LIST OF TABLES	xix
LIST OF FIGURES	xxi
LIST OF ABBRIVATIONS	xxv
LIST OF APPENDIX	xxviii
 CHAPTER	
 1 GENERAL INTRODUCTION	 1
1.1 Hypothesis	4
1.2 Objectives	5
 2 LITERATURE REVIEW	 6
2.1. Overview of <i>Plutella xylostella</i> L.	6
2.1.1. Systematic	6
2.1.2. Morphology	7
2.1.3. Biology	7
2.1.4. Host Plant	8
2.1.5. Distribution	9
2.1.6. Migration	11
2.1.7. Damages and economic loses	11
2.2. Management of <i>P. xylostella</i>	12
2.2.1. Chemical Control	12
2.2.2. Cultural Control	13
2.2.3. Biological control of DBM	13
2.3. Baculoviruses	23
2.3.1. Historical Background	24
2.3.2. Baculovirus Taxonomy and Identification Studies	25
2.3.3. Pathology and Replication of Baculoviruses	31
2.3.4. Ecology and Environment	32
2.3.5. Baculovirus Production and Field Application	34
2.4. Baculoviruses as Biopesticide	34
2.4.1. Recombinant Baculovirus Biopesticides	38
2.4.2. Formulation of baculoviruses	39
 3 MORPHOLOGICAL CHARACTERISTICS OF <i>P. XYLOSTELLAL</i>. GRANULOVIRUS AND CONFIRMATION OF TAIWANESE ISOLATE OF PXGV	 49

3.1.	Introduction	49
3.2.	Materials and Methods	51
3.2.1.	Insect Larvae	51
3.2.2.	Rearing of <i>P. xylostella</i> on Artificial Diet	51
3.2.3.	Artificial diet	52
3.2.4.	Source of Virus	55
3.2.5.	<i>In vivo</i> propagation of <i>P. xylostella</i> granulovirus	55
3.2.6.	Purification of <i>P. xylostella</i> GV	57
3.2.7.	DNA Extraction and Restriction Endonuclease (REN) Analysis	59
3.2.8.	Electron Microscopy Negative Staining	60
3.3.	Results	60
3.3.1.	Confirmation of Taiwanese Isolate	60
3.3.2.	Morphological Characteristics of PxGV	62
3.3.3.	Characteristics of Infected Larvae	64
3.4.	Discussion	66
3.5.	Conclusion	68
4	EFFECT OF VIRUS CONCENTRATIONS OF <i>P. XYLOSTELLA</i> GRANULOVIRUS ON DIFFERENT LARVAL INSTARS OF DIAMONDBACK MOTH	69
4.1.	Introduction	69
4.2.	Materials and Methods	70
4.2.1.	Source of Larvae and virus	70
4.2.2.	Virus Propagation and Purification	70
4.2.3.	Counting of Occlusion Bodies (OB _s)	71
4.2.4.	Virus Concentrations on Different Larval Instars and Estimate Median Lethal Concentration (LC ₅₀) and Median Lethal Time (LT ₅₀)	73
4.2.5.	Bioassay Tests	73
4.2.6.	Experimental Design and Statistics	75
4.3.	Results	76
4.3.1.	Influence of Larval Instars and Viral Concentrations on Larval Mortality	76
4.3.2.	Median Lethal Concentration (LC ₅₀) and Median Lethal Time (LT ₅₀) values for PxGV	81
4.4.	Discussion	83
4.5.	Conclusion	86
5	EFFICAY OF <i>P. XYLOSTELLA</i> GRANULOVIRUS (PxGV) TO ULTRAVIOLET RADIATION AND EFFECTS OF ADJUVANTS ON INFECTIVITY OF PxGV	87
5.1.	Introduction	87
5.2.	Materials and Methods	89
5.2.1.	Source of Larvae and Virus Propagation and Purification	89
5.2.2.	Virus Susceptibility to Ultraviolet	89
5.2.3.	Radiation Source and Exposure Condition	90
5.2.4.	Ultraviolet Protectants and Additives	91
5.2.5.	Effect of Ultraviolet Protectants and Additives on PxGV Infectivity	91

5.2.6.	Effect of Ultraviolet Protectants and Additives on <i>PxGV</i> Pathogenecity before Exposed to UV Radiation	91
5.2.7.	Bioassay Tests	92
5.2.8.	Experimental Design and Data Analysis	92
5.3.	Results	94
5.3.1.	Virus Susceptibility to Ultraviolet Radiation	94
5.3.2.	Effect of UV Protectants and Additives on <i>PxGV</i> Infectivity	98
5.3.3.	Effect of UV Protectants and Additives on <i>PxGV</i> Infectivity before Exposure to UV-B Radiation	101
5.3.4.	Median Lethal Concentration (LC ₅₀) of DBM larvae treated with <i>PxGV</i> + Adjuvants before Exposure to UV Radiation treated on DBM Second Larval Instar	104
5.3.5.	Median Lethal Time (LT ₅₀) of DBM larvae treated with <i>PxGV</i> + Adjuvants at Different Virus Concentrations before Exposure to UV Radiation	105
5.3.6.	Median Lethal Time (LT ₅₀) of <i>PxGV</i> + Adjuvants after 30 and 120 Minutes Exposure to UV Radiation	107
5.3.7.	Cumulative Mortality of DBM treated <i>PxGV</i> with Different Adjuvants after 30 and 120 Minutes Exposure to UV Radiation	108
5.4.	Discussion	110
5.5.	Conclusion	116
6	EFFECT OF SPRAY- DRIED FORMULATION ON THE INFECTIVITY OF <i>P. XYLOSTELLA</i> GRANULOVIRUS	118
6.1.	Introduction	118
6.2.	Material and Methods	120
6.2.1.	Preparation of Required Virus Concentration	120
6.2.2.	Preparation of Formulation	121
6.2.3.	Encapsulation Formulation Process	122
6.2.4.	Bioassay Tests	123
6.2.5.	Comparing of Different Formulations of Spray Draying before and after Exposure to UV-B Lamp	124
6.2.6.	Scanning Electron Microscopy and Particle Size Measurements	125
6.2.7.	Evaluation Suspensibility of Spray Dried Formulations	125
6.2.8.	Data Analysis	126
6.3.	Results	127
6.3.1.	Spary-dried Formulations	127
6.3.2.	Effect of Formulations after Spray Draying and Before Exposure to UV Radiation on Larval Mortality	129
6.3.3.	Effect of Spray-dried Formulations after Exposure to UV Radiation on Larval Mortality	130
6.3.4.	Median Lethal Concentration (LC ₅₀) and Median Lethal Time (LT ₅₀) of DBM Treated with CLM and KLM Spray-Dried Formulations	133
6.3.5.	Scanning Electron Microscopy and Particle Size Analysis	136
6.3.6.	Suspensibility of Spray-dried Formulations	138
6.4.	Discussion	139
6.5.	Conclusion	144

7	EVALUATION OF SPRAY-DRIED FORMULATIONS OF <i>P. XYLOSTELLA</i> GRANULOVIRUS IN A SEMI-FIELD TEST	146
7.1.	Introduction	146
7.2.	Materials and Methods	147
7.2.1.	Material Used for Semi-Field Test	147
7.2.2.	Application and Sampling	148
7.2.3.	Experimental Design and Data Analysis	151
7.3.	Results	151
7.4.	Discussion	157
7.5.	Conclusion	160
8	SUMMARY, GENERAL CONCLUSION AND RECOMMENDATION FOR FUTURE RESEARCH	162
8.1.	Summary	162
8.2.	General Conclusion	168
8.3.	Recommendation for Future Research	169
	REFERENCES	171
	APPENDICES	190
	BIO DATA OF STUDENT	201
	LIST OF PUBLICATIONS	202